

DEVELOPING “ALL NATURAL” PERSONAL CARE PRODUCTS: AN OVERVIEW OF ANTIMICROBIAL AND PRESERVATIVE OPTIONS

Sabinsa Corporation

Authors:

Muhammed Majeed, Ph.D. & Lakshmi Prakash, Ph.D.

info@sabinsa.com

www.sabinsa.com

www.sabinsacosmetics.com





Introduction

The need to control microbial activity is inherent in two major facets of personal care, namely, antimicrobial care wherein the product helps to retard the growth of topical pathogens or eliminates them, and preservatives that retard the growth of microbes in personal care product formulations.

In recent years, there has been increased dialogue related to natural antimicrobials as topical actives and preservatives in the personal care industry. Synthetic compounds long accepted as effective in controlling microbial growth have come under scientific and regulatory scrutiny. These efforts are mainly driven by safety and environmental concerns, and the increased incidence of antibiotic resistant microbial strains. Natural alternatives derived from botanicals are therefore being explored by researchers across the world.

TOPICAL ANTIMICROBIALS: AN INTRODUCTION



Like all biological materials, the skin harbors resident microflora. Under certain conditions and in some groups of people, the skin microflora grow invasively and trigger pathological processes resulting in acne, infections and skin eruptions. Invasive growth of certain microorganisms causes scalp infections and infections in the oral cavity, as well. Certain microorganisms found in the skin, scalp and nails may also generate malodor and irritation, affecting general health and well being. Antimicrobials find functional applications in cleansing formulations, deodorants and topical protectant formulations.

Examples of skin pathogens include *Propionibacterium acnes* a microorganism associated with acne and other skin infections, *Staphylococcus aureus* a bacterial strain found in infected wounds and skin eruptions including acne, *Staphylococcus epidermidis* a bacterial strain occurring in a variety of opportunistic bacterial skin infections and in acne, corynebacteria and micrococci that cause body odor. Athlete's foot (*Tinea pedis*) and onychomycosis (nail fungus) are other infections associated with the skin and nails. Examples of oral cavity pathogens include *Streptococcus mutans*, a bacterial strain associated with the progression of dental caries, Examples of scalp pathogens include *Pitysporum ovale* (*Malassezia furfur*, yeast strain associated with dandruff).

Microorganisms also affect dental health. Gum disease involves bacterial growth and production of metabolic substances that gradually destroy the tissue surrounding and supporting the teeth. These bacteria grow and attack the tissues causing "gingivitis" characterized by inflamed gums that bleed easily. If left untreated the condition progresses to "periodontal disease" wherein there is severe inflammation, bone damage and tooth loss. The causative bacteria reside in plaque, a deposit that forms on the base of the teeth and hardens to form "tartar". Poor oral hygiene is the major cause of gum disease. Lifestyle, nutrition and the ageing affect the immune response, and increase the risk of gum disease. Antimicrobials target oral pathogens such as *Streptococcus mutans*, while anti-inflammatory and wound healing extracts offer support to healthy gums and teeth.



Antibiotic resistant strains of microorganisms arise from non-judicious use of conventional antibiotics. There is therefore a need for natural products that effectively inhibit the growth of microorganisms. Multifunctionality is an additional advantage of natural extracts. Several of them offer anti-inflammatory, immunological and wound healing support as well.

In the last few years, a number of laboratory studies have revealed the efficacy of plant extracts and phytochemicals as antimicrobials. These properties are attributed to the presence of secondary metabolites such as phenolics in essential oils and tannins in herbal extracts. Some examples of classes of natural materials that afford antimicrobial protection include essential oils such as tea tree oil, rosemary oil, turmeric oil; plant extracts such as rosemary extract, sage extract, lemon balm extract, green tea extract, *Kaempferia galanga* extract, Neem leaf extract and oil, and isolated phytochemicals such as cinnamates, benzoates, eugenol.

NATURAL ANTIMICROBIALS: ILLUSTRATIVE EXAMPLES

COLEUS OIL (INCI:COLEUS FORSKOHLII ROOT OIL)



Coleus oil is an essential oil extracted from the roots of *Coleus forskohlii* a plant from the Natural Order Labiatae (Lamiaceae), a family of mints and lavenders. This species is a perennial herb with fleshy, fibrous roots that grows wild in the warm sub-tropical temperate areas in South Asia. The roots are eaten as a condiment or pickle in India. In recent years *Coleus forskohlii* has gained pharmacological importance as the only known plant source of the biologically active compound, forskolin,¹ an coleus oil is an useful byproduct of forskolin extraction. The newly discovered antimicrobial properties of the oil (of specific composition obtained using a proprietary extraction process)², render it useful in topical preparations.

Compounds such as 3-decanone (about 7%), bornyl acetate (about 15%), sesquiterpene hydrocarbons and sesquiterpene alcohols in major concentrations imparts unique pleasing spicy notes to the essential oil. β -sesquiphellandrene (about 13%) and γ -eudesmol (12.5%) were identified in experimental studies on the oil.³



In laboratory studies, Coleus oil was found to more effectively inhibit the growth of skin pathogens including *Propionibacterium acnes* (associated with acne), *Staphylococcus aureus* (a bacterial strain found in infected wounds and skin eruptions including acne), *Staphylococcus epidermidis* a bacterial strain occurring in a variety of opportunistic bacterial skin infections and in acne, than the well known tea tree oil. Additionally, Coleus oil was found to inhibit the yeast culture *Candida albicans* more effectively than tea tree oil.² The extract is safe to use in cosmetic formulations, it does not irritate the skin and its pleasant woody aroma blends with cosmetics.

KAEMPFERIA GALANGA (INCI: KAEMPFERIA GALANGA ROOT EXTRACT)

Kaempferia galanga (Lesser galangal, kencur), and *Alpinia galanga* (greater galangal) from the Zingiberaceae family are commonly used as spice ingredients and medicinal herbs in South-east Asia and are valued traditionally for their protective topical effects when applied as pastes.



Kaempferia galanga rhizome contains about 1.5 to 2% essential oil, whose main components are ethyl cinnamate (25%), ethyl-p-methoxycinnamate (30%) and p-methoxycinnamic acid. *Kaempferia galanga* is a good natural source of a biologically active ester compound ethyl p-methoxycinnamate.⁴ The antifungal action of ethyl-p-methoxycinnamate is reported in literature.⁵

In laboratory studies, the fraction of the extract rich in ethyl p-methoxycinnamate was found to effectively inhibit the growth of *Propionibacterium acnes*, producing significant zones of inhibition at concentrations even as low as 0.5%.⁶

TURMERIC OIL (INCI: CURCUMA LONGA) CURCUMA LONGA ROOTS, THE SOURCE OF TURMERIC OIL

Turmeric oil is obtained by steam distillation or solvent extraction of the powdered rhizome of species of the genus *Curcuma* (Family: Zingiberaceae). Of these species, *Curcuma longa* is the most well known. *Curcuma longa* yields 0.3-7.2% (usually 4-5%) of turmeric oil following steam distillation. The chief constituents of the essential oil are turmerone (60%) and related compounds, and zingiberene (25%).⁷



In India, turmeric is known for its cosmetic and wound healing properties since ancient times. The essential oil has been used as a perfume component and studies have shown that it furnishes antibacterial/antifungal⁸⁻¹¹, anti-inflammatory, insect-repellant activities and is effective in the treatment of scabies.

**OLEUROPEIN: INCI: OLEA EUROPAEA (OLIVE LEAF EXTRACT)
OLIVE LEAVES, THE SOURCE OF OLEUROPEIN**

Oleuropein is a polyphenolic compound found in plants belonging to the Oleaceae family, of which the olive tree is a member. The olive tree (*Olea europaea*) was known in biblical times as the “Tree of Life”.



Oleuropein is the major phenolic constituent extracted from olive leaf, (constituting about 19% (w/w)) and from the polar fraction of virgin olive oil. Oleuropein is reported to be hydrolyzed to another biologically active compound, hydroxytyrosol, *in vivo*. A variety of antimicrobial actions of oleuropein and its associated compounds have been demonstrated *in vitro*. The inhibitory action of oleuropein against the growth and toxin production of *Staphylococcus aureus*, *Bacillus cerus*, *Pseudomonas syringae* and several other bacterial strains (by oleuropein or its hydrolysis products) *in vitro*, is documented. For example, the presence of low concentrations (0.1% w/v) of oleuropein delayed the growth of *Staphylococcus aureus* in nutrient media, while higher concentrations (0.4-0.6% w/v) inhibited growth completely. Concentrations of oleuropein greater than 0.2% w/v inhibited growth and enterotoxin production in both types of media.

Oleuropein had bactericidal effects against a broad spectrum of gram-positive and gram-negative bacteria, but no effect was observed against yeast.¹² Antiviral properties¹³ are also reported *in vitro* and *in vivo*. Although the precise mechanism of antimicrobial action has as yet to be elucidated,^{12,14} oleuropein and related compounds appear to have surface-active properties that interfere with microbial cell membranes. Oleuropein could also interfere with the synthesis of amino acids that are crucial to viral replication and in the case of retroviruses, neutralize the production of reverse transcriptase and protease. Additionally, oleuropein is also reported to stimulate phagocytosis or the immune response to infection by pathogens.¹²



NEEM OIL AND NEEM LEAVES EXTRACT (INCI: MELIA AZADIRACHTA)



The Neem tree is traditionally labeled as “The Village Pharmacy” on account of its multifaceted healthful properties. Its properties range from immunomodulatory and anti-inflammatory effects to antimicrobial and pesticidal attributes. The leaves and seeds of Neem yield limonoids with wide biological applications. These have antibacterial, antiviral, insect repellent, anti-protozoal and anti-helmenthic properties.

The seed oil at concentration of 0.3% on agar plates were active against *Stahylococcus aureus* and at 0.4% was active against *Salmonella typhosa*. The seed oil at concentration of 3% on agar plate, was active against *E.coli* and *Proteus* species, a concentration of 6.0% was active against *Klebsiella pneumoniae*. Methanol and Butyl-methyl-ether extracts showed antifungal activity against the following strains of fungi including *Epidermaphyton floccosum*, *Microsporum canis*, *Microsporum gypseum*, *Trichophyton concentricum*, *Trichophyton entagrophytes*, *Trichophyton rubrum* and *Trichophyton violaceum*.¹⁵

A recent study determined the efficacy of a mucoadhesive dental gel containing *Azadirachta indica* leaf extract (25 mg/g). using commercially available chlorhexidine gluconate (0.2% w/v) mouthwash as positive control. The results of the study suggested that the dental gel containing neem extract significantly reduced plaque index and bacterial count as compared to the control group.¹⁶

GREEN TEA EXTRACT: (INCI: CAMELLIA SINENSIS LEAF EXTRACT)


The catechins in green tea were found to inhibit *Staphylococci* and *Yersinia enterocolitica*. Green tea extracts may make strains of drug-resistant bacteria more sensitive to penicillin. *In vitro* studies on particular antibiotic resistant strains of *Staphylococcus aureus* revealed that addition of green tea extract induced reversal of penicillin resistance. It was found that epicatechin gallate, markedly lowered the minimum inhibitory concentration (MIC) of oxacillin and other beta-lactams.^{17,18} Extracts of green tea were found to strongly inhibit *Escherichia coli*, *Streptococcus salivarius* and *Streptococcus mutans*, microorganisms found in the saliva and teeth of people suffering from dental caries.¹⁹ Green tea in combination with the synthetic antioxidant



butylated hydroxyanisole (BHA) reduced the hydrophobicity of *S. mutans* and greatly inhibited ($p < 0.001$) the formation of hyphae in *Candida albicans*. The increased antimicrobial activity of green tea is related to an impairment of the barrier function in microorganisms and a depletion of thiol groups.²⁰

NATURAL PRESERVATIVES

A preservative is essentially a chemical agent that will destroy microorganisms or prevent their growth in finished formulations. There is ample scope for microbial contamination at the end user level. It is therefore important that the cosmetic product be protected by a suitable preservative system to ensure safety in use and adequate shelf life.



Factors such as water activity and pH of the formulation, and the solubility and stability of the preservative in the base composition are factors that determine preservative efficacy. Additives such as glycerol, butylene glycol, salt (sodium chloride), sugars, soluble starches, dextrin, xanthan gum and others absorb water thereby lowering the water activity of the composition. Parallels in the list of natural actives include glucans, mannans and other polysaccharides extracted from commonly used culinary materials such as tamarind seed and fenugreek.

In the area of chemical preservatives that stop or inhibit microbial growth, common classes of chemicals used include acids, aromatic alcohols, N-methylol containing compounds, halogenated compounds, isothiazolinones, quaternary nitrogen compounds, and 1,2 diols. A number of phytochemicals found in natural sources resemble these compounds in chemical structure and are natural preservatives. These classes of phytochemicals include essential oil constituents (such as eugenol, thymol, carvacrol, terpenoid compounds), flavonoids, phenolic compounds, tannins and alkaloids. Some of these extracts when combined with ineffective antibiotics were found to be effective against antibiotic resistant strains.



Although no single natural extract has been found to be comparable in efficacy to conventionally used preservatives such as parabens, combinations of naturals with synergistic activity have been identified. Such extracts could offer protection against invasive growth of bacteria and fungi. Parabens for example, are poorly water soluble and their action is strongest against gram positive bacteria and fungi and weakest against gram negative bacteria.



Combinations of synthetic preservatives are therefore used in formulations and the safety and global regulatory acceptance of effective levels of such combinations is sometimes a problem. It is here that combinations of natural extracts may be useful. Very often such extracts also offer antioxidant action, with beneficial effects on product shelf life. One example is a proprietary natural extract of yellow curcuminoids from *Curcuma longa* (turmeric) roots³ and a colorless derivative Tetrahydrocurcuminoids (INCI:Tetrahydrodifuruloylmethane, Tetrahydrodemethoxydiferuloylmethane, Tetrahydrobisdemethoxydiferuloylmethane), that were found to offer more effective antioxidant protection than the conventionally used synthetic antioxidant butylated hydroxytoluene (BHT).²¹

Another example is resveratrol (Resvenox^{®4}) which offers antimicrobial and antioxidant benefits. Resveratrol is found in widely varying amounts in grapes (primarily the skins), raspberries, mulberries, blue berries, bilberries, cranberries, peanuts, scots pine, eastern white pine, and the roots and stalks of giant knotweed and Japanese knotweed (*Polygonum cuspidatum*). It has antibacterial and antifungal properties.²²



³ US Patent # 5,861,415, Sabinsa Corporation

⁴ A trademark of Sabinsa corporation



CONCLUSIONS

An overview of some of the natural alternatives to synthetic preservatives and antimicrobial functional ingredients in cosmetics is presented here. Synergistic blends of these materials would be functional, and simultaneously benefit the “all natural” image of personal care products. Please contact Sabinsa Corporation for further information on these and other natural ingredients for use in antimicrobial personal care, and as preservatives in finished formulations.



REFERENCES

1. De Souza, N.J. *Coleus forskohlii* Briq.- The Indian plant source for forskolin. Recent Advances in Medicinal, Aromatic & Spice crops, (ed: S..P. Raychaudhuri.) Today and Tomorrow's Printers and Publishers, New Delhi, India, 1991, Vol I: 83-91.
2. Misra, L.N. et al. Variability of the chemical composition of the essential oil of *Coleus forskohlii* genotypes. *J. Essential Oil Res.* 1994, 6:243-247.
3. United States Patent 6,607,712 Composition and methods containing an antimicrobial essential oil extended from *Coleus forskohlii*. Majeed, M., Prakash, S. (Aug. 19, 2003).
4. Kiuchi F, Nakamura N, Tsuda Y. 3-Carene-5-one from *Kaempferia galanga*. *Phytochemistry*, 26, 3350, 1987.
5. Gupta SK, Banerjee AB, Achari B Isolation of Ethyl p-methoxycinnamate, the major antifungal principle of *Curcuma zedoaria*. *Lloydia.* 1976 Jul-Aug;39(4):218-22.
6. Research Report (unpublished), Sabinsa Corporation, 2003.
7. Goldig, BT, Pombo, E, Samuel, CJ. Turmerones: Isolation from Turmeric and their Structure determination. *J. Chem. Soc. Chem. Commun.*1982, 363-364.
8. Chopra, R.N., Gupta, J.C., Chopra, G.S., (1971) *Ind. J. Med. Res.*, 29 (4), 769-772.
9. Ramprasad, C., Sirsi, M., (1956) *J. Sci. Ind. Res.*, 15c, 239-241.
10. Iyengar, M.A., Rao, M.P.R., Bairy, I., Kamath, M.S., (1984) *Indian Drugs* 31, 54-49.
11. Bhavani Shankar, T.N., Sreenivas Murthy, V. (1978) *J. Food. Sci. Tech.*, 15, 152-153.
12. Soler-Rivas, C. et al. (2000) Oleuropein and related compounds. *J. Sci. Food Agric.* 80:1013-1023.
13. Renis Harold. *In vitro* antiviral activity of calcium elnolate. In *Antimicrobial agents and chemotherapy*, 1969 (printed 1970) pp 167-279.
14. Juven B, Henis Y and Jacoby B, (1972). Studies on the mechanism of the antimicrobial action of oleuropein. *J Appl Bacteriol* 35:559-567.
15. Khan M, Schneider B, Wassilew and V Splanemann. The effect of raw material of neem tree, neem oils, neem extracts on dermatophytes, yeasts and molds. *Z. Hautkrankheiten* 63(6), 499-502, 1988.
16. Pai MR, Acharya LD, Udupa N. Evaluation of antiplaque activity of *Azadirachta indica* leaf extract gel--a 6-week clinical study. *J Ethnopharmacol.* 2004 Jan;90(1):99-103.
17. Yam TS et al. The effect of a component of tea (*Camellia sinensis*) on methicillin resistance, PBP2' synthesis, and beta-lactamase production in *Staphylococcus aureus*. *J Antimicrob Chemother* 1998 Aug;42(2):211-6,.
18. Shiota S et al. Marked reduction in the minimum inhibitory concentration (MIC) of beta-lactams in methicillin-resistant *Staphylococcus aureus* produced by epicatechin gallate, an ingredient of green tea. *Biol Pharm Bull* 1999. 22(12):1388-90.
19. Rasheed A and Haider M Antibacterial activity of *Camellia sinensis* extracts against dental caries. *Arch Pharm Res* 1998, 21(3):348-52.
20. Simonetti G, Simonetti N, Villa A. Increased microbicidal activity of green tea (*Camellia sinensis*) in combination with butylated hydroxyanisole. *J Chemother.* 2004 Apr;16(2):122-7.



21. Majeed, M, Badmaev, V Shivakumar, U and Rajendran, R *Curcuminoids Antioxidant Phytonutrients* NutriScience Publishers, Inc. Piscataway New Jersey. (1995).
22. Chan MM. Antimicrobial effect of resveratrol on dermatophytes and bacterial pathogens of the skin. *Biochem Pharmacol.* 2002 Jan 15;63(2):99-104.

